


New Records for Indian States: Andaman & Nicobar Islands

V. P. Prasad



How to cite:

Prasad V.P. 2008. New Records for Indian States: Andaman & Nicobar Islands. *Rheedea* 15(1): 61-62.

<https://dx.doi.org/10.22244/rheedea.2008.18.01.11>

Published in print: 31.08.2008

Published Online: 31.08.2008

Maharashtra

Juncus prismatocarpus R. Br. is the only species of the genus reported from Maharashtra (Sharma *et al.*, 1996). Now we report *J. maritimus* Lam. though it has already been reported from Gujarat (Cooke, 1958; Shah, 1978) and Karnataka States (Sharma *et al.* 1984).

Juncus maritimus Lam. Encycl. 3:264. 1789; Hook.f., Fl. Brit. India 6:395. 1892; Cooke, Fl. Pres. Bombay 3:308. 1958 (Repr.); Shah, Fl. Gujarat 691. 1978.

Flowering & Fruiting: October-December

Habit and Habitat: Caespitose herbs in the saline soil in abandoned sugarcane cultivated fields.

Distribution: India (Gujarat, Karnataka, Maharashtra), Afghanistan, West world of the Atlantic, North and South America and Australia.

Specimen Examined: India, Maharashtra, Sangli district, Padalwadi, 05.12.2006, Shimpale 1695; Yadav 3546 (Herbarium of Botany Department of Shivaji University, Kolhapur).

Note: The species is known for bioaccumulation of heavy metals such as Cd, Cu and Zn. It is used for phytoremediation of soils and estuarine ecosystems (Almeida *et al.*, 2004).

Acknowledgements

Authors are thankful to the Professor and Head, Department of Botany for providing facilities and Dr

Omprakash Kalame, Department of Journalism, Shivaji University, Kolhapur, for the help rendered during field work.

Literature Cited

- Almeida, C. Marisa, Mucha Ana P., Vasconcelos M. Teresa S. D. 2004. Influence of the sea rush *Juncus maritimus* on metal concentration and speciation in estuarine sediment colonized by the plant. *Environ. Sci. Technol.* 38: 312-318.
 Cook, C. D. K. 1996. *Aquatic and Wetland Plants of India*. Oxford University Press, Walton Street, Oxford.
 Cooke, T. 1958 (Repr. ed.). *The Flora of Presidency of Bombay* Vol. 3. Botanical Survey India, Calcutta.
 Shah, G. L. 1978. *Flora of Gujarat State*. Vol. 1 & 2. Sardar Patel University Press, Vallabh Vidyanagar, Gujarat.
 Sharma B. D., Singh, N. P., Raghavan, R. S. and U. R. Deshpande 1984. Flora of Karnataka Analysis. Botanical Survey of India, Culcutta.
 Sharma, B. D., Karthikeyan, S. & N. P. Singh (Eds) 1996. Flora of Maharashtra State: Monocotyledons. Botanical Survey of India, Calcutta.

Vinod B. Shimpale and S. R. Yadav¹

Department of Botany, The New College, Kolhapur 416 012, Maharashtra, India. E-mail: shimpale@yahoo.com

¹Department of Botany, Shivaji University, Kolhapur 416 004, Maharashtra, India. E-mail: sryadavdu@rediffmail.com

Andaman & Nicobar Islands

Sunil & Sivadasan (2000) reported *Kyllinga polyphylla* Willd. ex Kunth from Kerala as an addition to the flora of India. However, Karthikeyan *et al.* (1989) had included this species in their *Flora Indicae Enumeratio: Monocotyledonae* without specifying the locality. While revising the genus *Kyllinga* Rottb. for Flora of India project, one specimen collected from the Nicobar Islands was found at CAL. Therefore, at present, this species is distributed in the country in two widely separated geographical zones.

Kyllinga polyphylla Willd. ex Kunth, Enum. Pl.2:134.1837; Koyama, Gard. Bull. 30:163.1977& in Dassan. Fosb., Rev. Handb. Fl. Ceylon 5:247.1985; Karthikeyan *et al.*, Fl.Ind.

Enum. Monocot. 61.1989; Sunil & Sivadasan, *Rheeedea* 10:81.2000. *Kyllinga aromaticata* Ridl., Trans. Linn. Soc. London 2: 146.1884. *Cyperus aromaticus* (Ridl.) Mattf. & Kükenth. in Engl. Pflanzenr. IV. 20(101):581.1936; Koyama, Bot. Mag. (Tokyo) 83: 186.1970; Kern, Fl. Males. I, 7(3): 656.1974.

The species can be differentiated from the most allied *Kyllinga melanosperma* Nees by the presence of 6-8 leafy involucral bracts subtending the inflorescence. It is indigenous to E. Africa and was introduced in Fiji, Solomon Islands, Malesia and Sri Lanka (Kern, 1974). The plant is found near brackish water, close to coastal areas.

Specimens Examined: Andaman & Nicobar Islands, North Nicobar, Car Nicobar, Mus, 28-9-1976, N.G. Nair 4525, (CAL). Kerala, Alleppy dist., Aroor, 16-5-2001, B. Sreedevi s.n., (Herbarium, S.D. College, Alleppy); Ernakulam dist., Moothakunnam, 30-12-2007, V.P. Prasad, 192732 (BSI).

Acknowledgements

Author is thankful to the Director, Botanical Survey of India, Kolkata, Joint Directors, Botanical Survey of India, Pune and Central National Herbarium, Howrah for facilities; Dr S. Karthikeyan, Ex-Deputy Director, Botanical Survey of India, Pune for helpful suggestions; Dr M.S. Binojkumar, S.D. College, Alleppy, Kerala for sending a specimen on loan, and Dr. C.N. Sunil, S.N.M. College, Maliankara, Ernakulam for helping to locate the plant at Moothakunnam, Ernakulam district.

Literature Cited

- Karthikeyan, S., Jain, S. K., Nayar M. P. & M. Sanjappa 1989. *Florae Indicae Enumeratio: Monocotyledonae*. Botanical Survey of India, Kolkata. p. 61.
- Kern, J.H. 1974. Cyperaceae. In: C.G.G.J. van Steenis, *Flora Malesiana* Ser.1, Vol. 7. p. 656.
- Koyama, T. 1977. The Cyperaceae Tribe Cypereae of Ceylon. *Gard. Bull. Singapore* 30: 163.
- Koyama, T. 1985. Cyperaceae. In: M.D. Dassanayake & F.R. Fosberg, *A Revised Handbook to the Flora of Ceylon*. Amerind Publishing Co. Pvt. Ltd., New Delhi. Vol. 5. pp. 247-248.
- Sunil, C.N. & M. Sivadasan 2000. *Kyllinga polyphylla* Willd. ex Kunth (Cyperaceae): A new record for India. *Rheedia* 10: 81-83.

V.P. Prasad

Botanical Survey of India, Western Circle, Pune 411 001, Maharashtra , India. E-mail: Prasad_parur @ yahoo.com