




Notes on the Extended Distribution of Three Indian Endemics in Tamil Nadu

Ravikumar K., Lakshmanan V. & R. Vijaya Sankar



How to cite:

Ravikumar K., Lakshmanan V. & R. Vijaya Sankar 2004. Notes on the Extended Distribution of Three Indian Endemics in Tamil Nadu. *Rheedeia* 14(1&2): 69–70.

<https://dx.doi.org/10.22244/rheedeia.2004.14.1&2.10>

Received: 8.04.2003

Revised and accepted: 08.11.2004

Published in print: 31.12.2004

Published Online: 01.01.2022

Notes on the Extended Distribution of Three Indian Endemics in Tamil Nadu

Abstract

Diospyros angustifolia (Miq.) Kosterm., *Gomphostemma keralensis* K. Vivekananthan, R. Gopalan & R. Ansari and *Hedyotis bourdillonii* (Gamble) R.S. Rao & Hemadri are Indian endemics not reported so far from Tamil Nadu. Explorations in Theni district helped to locate these species from Pachakumatchi hills. Nomenclature and description of these endemics are provided.

Keywords: *Diospyros angustifolia* (Miq.) Kosterm., *Gomphostemma keralensis* K. Vivekananthan et al. and *Hedyotis bourdillonii* (Gamble) R.S. Rao & Hemadri, occurrence, Tamil Nadu

Introduction

Extensive field surveys conducted in Pachakumatchi Hills (High Wavy Mountains) in Theni district, Tamil Nadu under the Task Force Programme of the Botanical Survey of India enabled to locate *Diospyros angustifolia* (Miq.) Kosterm. (Ebenaceae), *Gomphostemma keralensis* K. Vivekananthan, R. Gopalan & R. Ansari (Lamiaceae) and *Hedyotis bourdillonii* (Gamble) R.S. Rao & Hemadri (Rubiaceae) as new additions to the flora of Tamil Nadu (Gamble, 1921; Matthew, 1983; Henry *et al.* 1987). They were the Indian endemics reportedly occurring outside the state boundary. (Saldanha & Ramesh, 1984; Mohanan *et al.* 1984). Brief descriptions of these species and notes on their distribution are given below.

Diospyros angustifolia (Miq.) Kosterm. in Ceylon J. Sci. 12: 106. 1977; Saldanha & Ramesh in Saldanha, Fl. Karnataka 1: 335. 1984. *Maba angustifolia* Miq., Anal. Bot. India 3: 13. 1852; Almeida, Fl. Maharashtra 3A: 178. 2001. *M. nigrescens* Dalzell in Dalzell & A. Gibson, Bombay Fl.: 142. 1861; Cooke, Fl. Bombay: 97. 1904; Gamble, Fl. Madras: 540. 1921. *Diospyros nigrescens* (Dalzell) Saldanha in Saldanha & Nicols., Fl. Hassan: 197. 1976; Kulkarni, Fl. Sindhudurg 240. 1988.

Trees, 4 to 8 m tall. Young branchlets densely brown-strigose. **Leaves** ovate-lanceolate to elliptic-lanceolate, 2-7 x 0.8-2.5 cm, coriaceous, narrowly acute at apex, subobtusate to cuneate at base, wavy, thickened and recurved along margin, ciliate when young. **Flowers** 3-merous, axillary. **Male flowers** 1-3 per cluster; calyx-teeth ovate, acute, hirsute, brown-strigose; corolla white, c. 5 mm long, lobes 3, obovate, obtuse, densely hairy without, stamens 9. **Female flowers** solitary, c. 5 mm across; calyx-lobes 3, ovate,

c. 5 x 5 mm, acute, sparsely strigose; corolla white, c. 5 mm long, lobes 3, oblong, obtuse, hairy without; ovary 3-celled; stigma 3-fid. **Berries** obliquely ellipsoid, c. 1.5 x 1.2 cm, rusty-brown when young with persistent style and calyx cup.

Flowering & Fruiting: May-August.

Habitat: Along streamlets in moist deciduous to evergreen forests, frequent.

Specimens Examined: **Tamil Nadu**, Theni District, Vellimalai Estate, along Sabarigiri river, *Lakshmanan* 95240 (MH). **Karnataka**, Hassan District, Shiradi Ghat, on the banks of Kempuhole, *Saldanha* 16560 (JCB).

Note: Canara in Karnataka is the type locality of this species (Type: *Metzius* 389, K). It is so far recorded only from Maharashtra and Karnataka. Kostermans (1981) stated that this species was close to the Sri Lankan endemic, *D. rheophytica* but differs in having dense strigose branchlets.

Gomphostemma keralensis K. Vivekananthan, R. Gopalan & R. Ansari in Kew Bull. 38: 189. 1983; Mohanan, Pandurangan & Raju in J. Econ. Tax. Bot. 5 (2): 455-459. 1984.

Subshrubs, to 1 m tall; stem 4-angled, stellate tomentose. **Leaves** elliptic-oblongate, 15-25 x 6-10 cm, chartaceous, glabrescent above, acuminate at both ends, dentate along margin; petioles 2-5.5 cm long. **Verticils** c. 2 cm across; bracts c. 1.5 x 0.5 cm. **Flowers** yellow, c. 1 cm long; calyx c. 1.2 cm long; corolla puberulous without. **Nutlets** oblong-ellipsoid, c. 6 x 3 mm, pubescent.

Flowering & Fruiting: June-September

Habitat: Grasslands adjacent to evergreen forests, rare.

Specimens Examined: **Tamil Nadu**, Theni district, Vellimalai, *Lakshmanan* 87928 (MH). **Kerala**, Idukki district, Kulamavu, *Mohanan* 74544 (MH).

Note: Mangaladevi, Periyar Wildlife Sanctuary in Idukki (Kerala) is the type locality of this species (*Vivekananthan* 48658, holotype, CAL; isotype MH). Later, it was located from Kulamau of the same district by Mohanan *et al.* (1984).

Hedyotis bourdillonii (Gamble) R.S. Rao & Hemadri in Indian For. 99: 378. 1973; N.C. Nair, V. J. Nair & Ansari in Bull. Bot. Surv. India 22: 205. 1980. *Oldenlandia bourdillonii* Gamble in Bull. Misc. Inform. 1919. 404. 1919 & Fl. Madras 422. 1921. *Hedyotis silent-valleyensis* Vajravelu, N.C. Rathakrishnan & P. Bhargavan, J. Bombay Nat. Hist. Soc. 80 (2): 402. 1983; Deb & Dutta in J. Bombay Nat. Hist. Soc. 83 (1): 466. 1986.

Shrubs, 1-1.5 m tall, branchlets dichotomous, hirsute. **Leaves** elliptic-lanceolate, 3-6 x 1.5-2.5 cm, membranous, hirsute, obtuse at base, acuminate at apex; lateral nerves 3-4 pairs, oblique; stipules triangular, 1.5-2.5 x 2.5-4.5 mm,

pubescent; teeth 9-11, pectinate, villous. *Inflorescence* a terminal dichasial cyme. *Flowers* subsessile, c. 7 mm long; bracts leafy, 6-8 mm long, villous; calyx 4 lobed, triangular, acute, pubescent, white to pale pink; corolla tubular, 7-8 mm long, pubescent without, 4 lobed, oblong, incurved at apex, serrulate along margin, pale blue to pale purple. *Capsules* oblong, c. 6 x 3 mm, densely villous at base; seeds 8-10, narrowly 3 - winged, reticulate, pale brown.

Flowering & Fruiting: December-February and June-August.

Habitat: Grasslands at high altitudes, locally common.

Specimens Examined: **Tamil Nadu**, Theni district, Pachakumatchi hills, Maharajamettu peak and Nursery Valley, Lakshmanan 87536, 95253 (MH). **Kerala**, Palaghat district, Silent Valley, Ansari 51466 (MH).

Note: Gamble (1919) described this species based on Bourdillon's collection in 1887 from the Travancore hills in Kerala after T. F. Bourdillon 111(K) and T.F. Bourdillon s.n. (MH Accn. No. 24321). Nair *et al.* (1980) relocated it from Silent Valley after 120 years of the type collection.

Acknowledgements

The authors are thankful to Dr P. Daniel, (Retd) Joint Director, Botanical Survey of India, Southern Circle, Coimbatore for critical comments on the manuscript and to Shri. D. K. Ved, Additional Director, FRLHT, Bangalore for facilities and encouragement.

Literature Cited

- Deb, D.B. & R. Dutta 1986. On the identity of *Hedyotis silent-valleyensis* (Rubiaceae). *J. Bombay Nat. Hist. Soc.* 83(1): 466.
- Gamble, J.S. 1919. *Flora of the Presidency of Madras* 2. Adlard & Sons Ltd., London.
- Henry, A.N., Kumari, G.R. & V.Chithra 1987. *Flora of Tamil Nadu, India* 1: Analysis 2. Botanical Survey of India, Coimbatore.
- Kostermans, A.J.G.H. 1981. Ebenaceae. In: Dassanayake, M.D. and F.R. Fosberg (Eds), *A Revised Handbook to the Flora of Ceylon* 3. Oxford & IBH Publishing Co. Pvt. Ltd., New Delhi. p. 9.
- Matthew, K. M. 1983. *The flora of the Tamil Nadu Carnatic*. 2. The Rapinat Herbarium, St. Joseph's College, Tiruchirapalli.
- Mohanan, C.N., Pandurangan, A.G. & V.S. Raju 1984. Some rare and interesting angiosperm taxa from the forests of Idukki Hydro-electric project area, Kerala, India. *J. Econ. Tax. Bot.* 5(2): 455-459.
- Nair, N.C., Nair, V. J. & R. Ansari 1980. Notes on some rare plants from South India. *Bull. Bot. Surv. India* 22(1-4): 205-207.
- Saldanha C. J. & B. R. Ramesh 1984. Ebenaceae. In: Saldanha, C. J. *Flora of Karnataka* 1. pp.334-341. Oxford & IBH Publishing Co., New Delhi.

Received 8.4.2003

Revised and accepted 8.11.2004

K. Ravikumar, V. Lakshmanan¹ and R. Vijaya Sankar

Foundation for Revitalisation of Local Health Traditions, # 74/2, Jarakabande Kaval, Attur P.O., Via Yelahanka, Bangalore 560 064, Karnataka, India. e-mail: k.ravikumar@ftrlht.org.in

¹Botanical Survey of India, Southern Circle, TNAU Campus, Coimbatore 641 003, Tamil Nadu, India.